

Pion properties at finite density

Seung-il Nam^{1,*} and Hyun-Chul Kim^{2,†}

¹*Yukawa Institute for Theoretical Physics, Kyoto University,*

Kitashirakawaoiwake, Sakyo, Kyoto 606-8502, Japan

²*Department of Physics, Inha University,*

253 Yonghuyn-dong, Nam-gu Incheon 402-751, Korea

(Dated: August 28, 2008)

Abstract

In this talk, we report our recent work on the pion weak decay constant (F_π) and pion mass (m_π) using the nonlocal chiral quark model with the finite quark-number chemical potential (μ) taken into account. Considering the breakdown of Lorentz invariance at finite density, the time and space components are computed separately, and the corresponding results turn out to be: $F_\pi^t = 82.96$ MeV and $F_\pi^s = 80.29$ MeV at $\mu_c \approx 320$ MeV, respectively. Using the in-medium Gell-Mann-Oakes-Renner (GOR) relation, we show that the pion mass increases by about 15% at μ_c .

PACS numbers: 12.38.Lg, 13.20.Cz, 14.40.Aq.

Keywords: pion weak decay constant, pion mass, finite density, instanton vacuum.

*E-mail: sinam@yukawa.kyoto-u.ac.jp

†E-mail: hchkim@inha.ac.kr

I. INTRODUCTION

The pion is identified as the Nambu-Goldstone (NG) boson from the spontaneous breakdown of chiral symmetry (SB χ S) which is essential in describing low-energy hadronic phenomena. Thus, the in-medium modification of the F_π and m_π is of great importance to understand the chiral symmetry restoration in matter. Experimentally, the modifications of these quantities can be measured from deeply bound pionic atoms.

There has been a great amount of theoretical works on this subject. For example, meson-baryon chiral perturbation theory (χ PT) and models with chiral symmetry were applied for this purpose [1, 2, 3]. Since the Lorentz invariance is broken in medium, one has to study the space and time components of the pion weak decay constant separately. In in-medium χ PT [4], the magnitude of its space component F_π^s was shown to be about four times smaller than that of the time component F_π^t at ρ_0 . In the QCD sum rules, it was discussed that dimension-five operators are responsible for making splitting between F_π^t and F_π^s , and the contributions of the intermediate Δ state makes F_π^s much smaller than F_π^t [3].

In the present work, we investigate the modifications of the F_π and m_π for the finite quark-number chemical potential ($\mu \neq 0$) but at zero temperature ($T = 0$), employing the nonlocal chiral quark model (NL χ QM), derived from the nontrivial instanton vacuum [5, 6]. We will show that $F_\pi^t = 82.96$ MeV and $F_\pi^s = 80.29$ MeV at the critical value $\mu_c \approx 320$ MeV, which are about $13 \sim 16\%$ smaller than that in free space ($F_\pi = 93$ MeV). The results are compatible with those obtained in other models. Using the GOR relation, satisfied within the model [5], we estimate the pion mass shift and find that the mass is increased by about 15 % at μ_c .

II. NONLOCAL CHIRAL QUARK MODEL WITH FINITE μ

The quark zero-mode solution in the presence of the instanton can be obtained with the finite μ as follows [6]:

$$[i\cancel{\partial} - i\gamma_4\mu - \cancel{A}_{I\bar{I}}] \Psi_{I\bar{I}}^{(0)} = 0. \quad (1)$$

A quark propagator deduced from Eq. (1) reads:

$$S = \frac{1}{i\cancel{\partial} - i\gamma_4\mu + iM(i\partial, \mu)}, \quad (2)$$

where M denotes the momentum-dependent and μ -dependent quark mass that arises from the Fourier transform of the quark zero-mode solution:

$$M(\bar{k}) = M_0(\mu)\bar{k}^2\psi^2(\bar{k}). \quad (3)$$

Here, $\bar{k} = (\vec{k}, k_4 + i\mu)$. The M_0 is the constituent quark mass at $k^2 = 0$, which depends on μ and becomes zero at the critical value $\mu_c \approx 320$ MeV, indicating the first-order phase transition [7]. The analytical expressions for ψ_4 and $\vec{\psi}$ are given in Ref. 6. Since the Lorentz invariance is broken for the finite μ , the PCAC relation in medium should be decomposed into the space (s) and time (t) parts as follows:

$$\langle 0 | \mathbf{A}^a(x) | \pi^b(P) \rangle = i\sqrt{2}F_\pi^s \delta^{ab} \mathbf{P} e^{-iP \cdot x}, \quad \langle 0 | A_4^a(x) | \pi^b(P) \rangle = i\sqrt{2}F_\pi^t \delta^{ab} P_4 e^{-iP \cdot x}. \quad (4)$$

In order to compute the relevant matrix elements, we write the effective chiral action in the presence of an external axial-vector source $J_{5\mu}^a$:

$$\mathcal{S}_{\text{eff}}[\pi, \mu, J_{5\mu}^a] = -\text{Sp} \ln \left[i\bar{\partial} + \gamma_5 \gamma^\mu \frac{\tau^a}{2} J_{5\mu}^a + \sqrt{M(i\bar{\partial}, J_{5\mu}^a)} U_5 \sqrt{M(i\bar{\partial}, J_{5\mu}^a)} \right]. \quad (5)$$

From this effective chiral action and the in-medium PCAC relation, we can obtain the following compact expressions for the F_π for finite μ :

$$\begin{aligned} F_\pi^s(\mu) &\approx F_\pi^{\text{exp}} + \mu^2 \left[\frac{N_c}{F_\pi^{\text{exp}}} \int \frac{d^4k}{(2\pi)^4} \left(\frac{8k_4^2 \tilde{\mathcal{M}}' \tilde{\mathcal{M}}''}{k^2 + \mathcal{M}^2} - \frac{10k_4^2 \tilde{\mathcal{M}}'^2}{[k^2 + \mathcal{M}^2]^2} \right) \right], \\ F_\pi^t(\mu) &\approx F_\pi^s(\mu) + \mu^2 \left[\frac{N_c}{F_\pi^{\text{exp}}} \int \frac{d^4k}{(2\pi)^4} \frac{8k_4^2 \tilde{\mathcal{M}}'^2}{[k^2 + \mathcal{M}^2]^2} \right]. \end{aligned} \quad (6)$$

For more details, one can refer Ref. 7.

III. NUMERICAL RESULTS

We now present numerical results for the F_π for finite μ . In the left panel of Fig. 1, we show F_π^s and F_π^t as functions of μ . At $\mu = 0$ we find $F_\pi^s = F_\pi^t \approx 93$ MeV as it should be, and we obtain $F_\pi^t \approx 82.96$ MeV and $F_\pi^s \approx 80.29$ MeV at $\mu_c \approx 320$ MeV. When we examine the ratio $F^{(s,t)}/F_\pi^{\text{exp}}$, it must be unity at $\mu = 0$ and then it is getting smaller gradually as μ increases. At μ_c , it turns out that $F_\pi^t/F_\pi^{\text{exp}} \approx 0.89$ and $F_\pi^s/F_\pi^{\text{exp}} \approx 0.86$. From these observations, F_π^s/F_π^t is less than unity for the whole region of the NG phase ($\mu \leq \mu_c$), and the F_π is reduced by about $13 \sim 16\%$. We summarize the results in Table I. In the QCD

sum rule [3], it was discussed that the splitting between the time and space components is represented by the dimension-five condensates. Especially, $F_\pi^s/F_\pi^{\text{exp}}$ becomes much smaller, when the intermediate Δ state is considered ($0.78 \rightarrow 0.57$). The ratios were also studied in in-medium χ PT in the heavy baryon limit [4]: It was found that $F_\pi^t/F_\pi^{\text{exp}} \approx 0.90$, which is compatible with the present results, whereas the space component is given much smaller: $F_\pi^s/F_\pi^{\text{exp}} \approx 0.25$ at ρ_0 .

TABLE I: Numerical results for $F_\pi^{s,t}$ and m_π for finite μ .

	F_π^s	F_π^t	m_π
$\mu = 0$	93 MeV	93 MeV	139.33 MeV
$\mu_c \approx 320$ MeV	80.29 MeV	82.96 MeV	160.14 MeV
Modification	16% $\downarrow$	13% $\downarrow$	15% $\uparrow$

Now, we would like to investigate the change of the pion mass for finite μ . For this purpose, we use the results for $F_\pi^{s,t}$ computed previously and the in-medium GOR relation:

$$(m_\pi^* F_\pi^t)^2 = 2m_q \langle i q^\dagger q \rangle^*, \quad (7)$$

where m_q is the current quark mass, taken to be around 5 MeV. In the right panel of Fig. 1, we draw m_π as a function of μ . In free space, we observe $m_\pi = 139.33$ MeV which is in good agreement with the experimental value, whereas $m_\pi = 160.14$ MeV at the critical value $\mu_c \approx 320$ MeV. This observation tells us that the pion mass increases almost linearly and at μ_c it becomes about 15% heavier than that in free space. We summarize this result also in Table I.

IV. SUMMARY AND CONCLUSION

We have investigated the pion weak decay constant and pion mass for finite μ within the framework of the nonlocal chiral quark model in the presence of the finite quark-number chemical potential. For the numerical results, we obtained $F_\pi^t = 82.96$ MeV and $F_\pi^s = 80.29$ MeV at $\mu_c \approx 320$ MeV. Considering the in-medium Gell-Mann-Oakes-Renner relation, we also studied the pion mass modification for finite μ . The results are summarized in Table I. However, the splitting between the time and space components of F_π has turned out to be relatively small in comparison to those from in-medium chiral perturbation theory and the




FIG. 1: F_π^s and F_π^t (left), and m_π (right) as functions of the quark-number chemical potential μ up to $\mu = \mu_c \approx 320$ MeV.

QCD sum rule. As discussed in the previous section, the Δ -state contribution may cause this difference. A further study including meson-loop corrections is under progress.

Acknowledgments

The authors are grateful to the organizers for the 4th Asia-Pacific Conference on Few-Body Problems in Physics 2008 (APFB08), which was held during 19 ~ 23 August 2008, in Depok, Indonesia. They also thank T. Kunihiro and S. H. Lee for fruitful discussions. S.i.N. was partially supported by the Grant for Scientific Research (Priority Area No.17070002 and No.20028005) from the Ministry of Education, Culture, Science and Technology (MEXT) of Japan. The present work is also supported by Inha University Research Grant (INHA-37453).

-
- [1] M. Kirchbach and D. O. Riska, Nucl. Phys. A **578**, 511 (1994).
 - [2] A. Wirzba and V. Thorsson, arXiv:hep-ph/9502314.
 - [3] H. c. Kim and M. Oka, Nucl. Phys. A **720**, 386 (2003).
 - [4] M. Kirchbach and A. Wirzba, Nucl. Phys. A **616**, 648 (1997).
 - [5] D. Diakonov and V. Y. Petrov, Nucl. Phys. B **272**, 457 (1986).
 - [6] G. W. Carter and D. Diakonov, Phys. Rev. D **60**, 016004 (1999).
 - [7] S. i. Nam and H. -Ch. Kim, Phys. Lett. B **666**, 324 (2008)